

# REMEDY

A fungicide for the control of Powdery Mildew and other diseases on terrestrial and indoor ornamental plants, greenhouse and garden crops and turf.



ACTIVE INGREDIENT:	
Potassium Bicarbonate .....	85.00%
OTHER INGREDIENTS .....	15.00%
TOTAL .....	100.00%
EPA Est. No. 4-NY -1	EPA Reg No. 70870-1-4

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

(See Back Panel for the First Aid Information)

60490203

TRUSTED SINCE 1926

**BONIDE**®

#### CONTROLS:

Blackspot, Downy Mildew, Powdery Mildew, Botrytis Blight and gray mold, Scab and other listed plant diseases.

#### Use On:

- Roses & Flowers
- Vegetable Crops
- Ornamental Nut and Fruit Trees as listed on label

Trusted since 1926  
**BONIDE PRODUCTS, INC.**

**Complete directions for use have been removed!**

**KEEP OUT OF REACH OF CHILDREN**

Dispose of properly  
Buyers Guarantee limited to label claims

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.

Questions, Concerns or Comments?

Write to  
BONIDE PRODUCTS, INC.  
Consumer Services  
6301 Sutliff Road  
Oriskany, NY 13424

Lift & Peel Here →

## REMEDY

**CONTROLS** powdery mildew, black spot, alternaria blight, botrytis blight and gray mold, fusarium leaf spot, septoria leaf spot...**USE** on woody and ornamental shade trees, also aster, azalea, chrysanthemum, dogwood, English ivy, geranium, holly, juniper, privet, viburnum.

Store and transport in an upright position.

EPA Est. No. 4-NY-1 EPA Reg. No. 70870-1-4  
Buyers Guarantee limited to label claims.

Distributed by  
**BONIDE PRODUCTS, INC.**  
Oriskany, NY 13424

### PRECAUTIONARY STATEMENTS

#### HAZARD TO HUMANS AND DOMESTIC ANIMALS

**CAUTION** - Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

#### FIRST AID

- If Swallowed:**
- Call poison control center or doctor immediately for treatment advice.
  - Have person drink several glasses of water.
  - Do not induce vomiting.
  - Do not give anything by mouth to an unconscious person.
- If In Eyes:**
- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
  - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
  - Call a poison control center or doctor for treatment advice.

#### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-228-5635 Ext. 007 for emergency medical treatment information.

## Page: One

### ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container in a cool, dry place. Prevent exposure to moisture.

**CONTAINER DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.

### GENERAL INFORMATION

REMEDY is a broad spectrum, contact foliar fungicide that controls a variety of diseases on terrestrial and indoor woody and herbaceous ornamental, greenhouse and garden plants and turf.

REMEDY is applied with spray equipment using a sufficient volume of water to insure complete coverage of all stems and foliage.

**IMPORTANT:** Read the entire label before using REMEDY.

## Page: Two

### MIXING AND APPLICATION INSTRUCTIONS

REMEDY must be diluted with water before applying. Carefully measure and mix the stated amounts of product and water.

Mix 2 level tablespoons of REMEDY per gallon of water. Do not exceed this use rate. Mix the solution thoroughly before use. One gallon of this solution is sufficient to treat approximately 450 square feet. Apply the solution to all exposed surfaces of the plant. Use a sufficient spray volume to obtain complete coverage of all foliage and stems. Uniform and complete coverage is essential for the most effective results.

### DO NOT STORE UNUSED SPRAY SOLUTIONS.

Spray adjuvants are not required. Some tank mixes with other pesticide products and fertilizers may produce spray mixes that are incompatible. A "jar test" is recommended for unfamiliar tank mixes. As with any pesticide, the addition of spray adjuvants and other pesticides to the spray tank may cause phytotoxicity to the foliage and/or fruit susceptible crops and plants.

### Application Timing

Start application at first sign of disease. For best protection, repeat at one to two week intervals until conditions are no longer favorable for disease development. Shorten the interval during rainy weather or during periods of high relative humidity.

For appropriate crops, Remedy may also be applied post-harvest to remaining foliage to decrease overwintering inoculum.

### ORNAMENTAL APPLICATIONS:

#### PLANT TYPES

**Remedy is for:** Woody and ornamental shade trees, Woody and herbaceous ornamentals, Ornamental nut and fruit trees, Annual and perennial flowers, Foliage plants, Ground Covers and Bedding plants.

## Page: Three

## Page: Four

**FOR USE ON:**

- |                         |                   |                                |
|-------------------------|-------------------|--------------------------------|
| *Ageratum               | *Delphinium       | *Lipstick Vine (Aeschynanthus) |
| *Ajuga                  | *Dianthus         | *Marigold                      |
| *Almond-Ornamental      | *Dizygotheca      | *Monarda (Bee Balm)            |
| *Alyssum                | *Dogwood          | *Nectarine-Ornamental          |
| *Andromeda              | *English Ivy      | *Pachysandra                   |
| *Aphelandra             | *Episcia          | *Palm                          |
| *Apple-Ornamental       | *Euonymus         | *Pansy                         |
| *Apricot-Ornamental     | *Fern             | *Peach-Ornamental              |
| *Artemisia              | *Forsythia        | *Peperomia                     |
| *Aster                  | *Gazania          | *Periwinkle                    |
| *Azalea                 | *Geranium         | *Philodendron                  |
| *Boxwood                | *Gladiolus        | *Phlox                         |
| *Calendula              | *Gloxinia         | *Pilea                         |
| *Carnation              | *Gypsophila       | *Pine                          |
| *Centaurea montana      | *Hawthorn         | *Pittosporum                   |
| *Cherry-Ornamental      | *Heliopsis scabra | *Plum-Ornamental               |
| *Chrysanthemum          | *Hibiscus         | *Poinsettia                    |
| *Cineraria              | *Holly            | *Poppy                         |
| *Citrus                 | *Hoya             | *Pothos                        |
| *Coleus                 | *Hydrangea        | *Primrose                      |
| *Columbine              | *Impatiens        | *Privet                        |
| *Coral Bells (Heuchera) | *Iris             | *Protea                        |
| *Crape Myrtle           | *Juniper          | *Prune-Ornamental              |
| *Crassula               | *Kalanchoe        | *Pulmonaria                    |
| *Croton                 | *Lavender         |                                |
|                         | *Lemon balm       |                                |
|                         | *Lipstick balm    |                                |

\*NOT FOR THIS USE IN CALIFORNIA

- |                 |             |           |
|-----------------|-------------|-----------|
| *Pyracantha     | *Schefflera | *Verbena  |
| *Rhododendron   | *Snapdragon | *Viburnum |
| Rose            | *Statice    | *Violet   |
| *Rose-of-Sharon | *Tree Ivy   | *Zinnia   |

**DISEASES CONTROLLED**

- |                       |                            |
|-----------------------|----------------------------|
| Alternaria blight     | Entomosporium leaf spot    |
| Anthraxnose           | Fusarium                   |
| Ascochyta blight      | Helminthosporium leaf spot |
| Black Spot            | Ink spot                   |
| Botrytis blight       | Phomopsis blight           |
| Botrytis gray mold    | Powdery mildew             |
| Cercospora leaf spot  | Ovulinia sp.               |
| Corynespora leaf spot | Ramularia leaf spot        |
| Didymellina leaf spot | Ray blight                 |
| Diplodia tip blight   | Scab                       |
|                       | Septoria leaf spot         |

**GARDEN APPLICATIONS:**

**FOR USE ON:**

- |              |              |              |                 |
|--------------|--------------|--------------|-----------------|
| Acorn squash | *Avocado     | *Cabbage     | *Cherries       |
| *Almonds     | *Basil       | *Caneberries | *Chicory        |
| Apples       | *Blueberries | Cantaloupe   | *Citrus         |
| *Apricots    | *Broccoli    | *Cauliflower | *Crenshaw melon |

\*NOT FOR THIS USE IN CALIFORNIA

**Page: Five**

**Page: Six**

**Garden Applications continued**

- |                 |             |                 |
|-----------------|-------------|-----------------|
| Cucumber        | *Peaches    | *Snow peas      |
| *Dry Beans      | *Pears      | *Spinach        |
| *Eggplant       | *Peas       | Strawberries    |
| *Endive         | *Pecans     | Summer squash   |
| *Escarole       | *Peppers    | *Sweet Potatoes |
| Grapes          | *Pistachio  | *Thyme          |
| Honeydew        | *Plums      | *Tomatoes       |
| *Kale           | *Potatoes   | *Turnip         |
| *Lettuce        | *Prunes     | *Walnut         |
| *Lima Bean      | Pumpkin     | Watermelon      |
| *Mango          | *Radicchio  | *Winter squash  |
| Muskmelon       | *Rosemary   | Zucchini        |
| *Mustard greens | *Sage       |                 |
| *Okra           | *Snap beans |                 |
| *Oregano        |             |                 |

\*NOT FOR THIS USE IN CALIFORNIA

**DISEASES CONTROLLED**

- |                      |              |                |
|----------------------|--------------|----------------|
| Alternaria leaf spot | Downy Mildew | Phoma          |
| Anthraxnose          | Fusarium     | Phytophthora   |
| Botrytis             |              | Powdery mildew |

**TURF APPLICATION**

Remedy is for use in the following use sites: Residential lawns and other turf grass sites.

**FOR USE ON**

Bentgrass, Bermudagrass, Blue grass, Kentucky annual, Centipedegrass, fescue, fine tall, chewings, red Ryegrass, perennial, annual, St. Augustinegrass, Zoysiagrass.

**DISEASES CONTROLLED**

Alternaria leaf spot, Anthracnose, Ascochyta blight, Cercospora leaf spot, Downy mildew, Helminthosporium leaf spot, Powdery mildew, Septoria leaf spot.

**Notice:** Buyer assumes all responsibility for safety if use is not in accordance with label instructions.

**IMPORTANT NOTICE:**

**DISCLAIMER AND LIMITATION OF LIABILITY**

This product has been researched to provide necessary data to support its use on the ornamental and garden plants listed on the label. However, it is understood that tests may not have been carried out on all varieties or cultivars and under all growing conditions on all plants listed on the label. Always follow the label directions. Exercise care when using this product on a given variety until familiar with the results under your growing conditions.

**Page: Seven**

**Page: Eight**